

The Relationship between Body Composition with Blood Pressure and Sleep Quality in Male Dormitory Student at Allameh Tabataba'i University

Mehdi Kushkestanti

Mohsen Parvani*

Shiva Ebrahimpour Nosrani

Seyed Yousef Bathaezadeh

M.Sc. Student of Exercise Physiology, Faculty of Physical Education and Sport Sciences, Allameh Tabataba'i University, Tehran, Iran

Received: November 14, 2019; **Accepted:** January 14, 2020

doi:10.22054/nass.2019.10536

Abstract

Background and Purpose: Hypertension and poor sleep quality can increase the risk of cardiovascular diseases, diabetes, stroke, etc. Therefore, the purpose of this study was to investigate the relationship between body compositions with hypertension and sleep quality in male students living in Allameh Tabataba'i University dormitory. **Material and Method:** The present study was descriptive-correlational. The statistical population consisted of 170 male dormitory students of Allameh Tabataba'i University who were randomly selected. First, body composition indices and blood pressure of subjects were measured and recorded. Then, the Pittsburgh Questionnaire were completed by students (PSQI) to assess their sleep quality. Data were analyzed using SPSS 21 software at a significant level ($P < 0.05$). **Results:** The results showed that there was a positive and significant correlation between systolic ($r = 0.281$, $p < 0.01$) and diastolic ($r = 0.357$, $p < 0.01$) blood pressure with fat percentage. Also, body mass index was significantly correlated with systolic ($r = 0.297$, $p < 0.01$) and diastolic blood pressure ($r = 0.322$, $p < 0.01$) while there was an inverse and significant correlation between muscle mass with systolic ($r = -0.252$, $p < 0.01$) and diastolic ($r = -0.356$, $p < 0.01$) blood pressure. Also, there was no significant relationship between students' sleep quality with fat mass ($r = -0.023$, $p = 0.76$) and muscle mass ($r = -0.015$, $p < 0.84$). **Conclusions:** In conclusion, according to the direct relationship between fat percentage and hypertension among students living in the dormitory, it can be concluded that the use of exercise in leisure time and recreational programs to improve body composition can play an important role in the prevention and treatment of hypertension in their future.

Keywords: Blood pressure, Sleep Quality, Muscle mass, Body fat percentage, Exercise

* **Author's e-mails:** mehdi.kushk@gmail.com, mparva2020@gmail.com
(**Corresponding author**), ebrahimpourshiva94@gmail.com,
yousef.bathae@gmail.com

2009; Suen, Ellis Hon, & Tam, 2008); because, it has been proven that sleep disorders influence on human health by changing the rhythm of hormone release, developing cardiovascular dysfunction, dyslipidemia, metabolic syndrome, and a poor blood glucose regulation (Bixler, 2009).

Control and identification of factors influencing high blood pressure and lower sleep quality are of great importance to improve the health status of the community. Assessment of body composition and anthropometric indices are of the simplest and most accessible methods that do not need expensive laboratory equipment. Several studies have examined the relationship between body composition and various disease-risk factors. Moreover, studies have reported that there is a positive correlation between hypertension with body mass index and fat percentage (Deng et al., 2012) while there is a negative correlation between hypertension and muscle mass (Ye et al., 2018). Furthermore, researchers have examined the relationship between BMI and sleep quality. In fact, there is a significant negative correlation between BMI and sleep quality (Vargas, Flores, & Robles, 2014). Also, many studies have reported that there is a significant negative correlation between subcutaneous fat percentage and visceral fat with sleep quality while there is a positive correlation between muscle mass and sleep quality (Rahe, Czira, Teismann, & Berger, 2015). (Kahlhöfer, Karschin, Breusing, & Bosy-Westphal, 2016).

Given the different lifestyles of dorm students and their vulnerability to peers, as well as the necessity for a healthy lifestyle to prevent different diseases throughout life, assessing risk factors such as body composition, sleep quality and blood pressure are of great importance. Over the past years, less research has examined the relationship between body composition with high blood pressure and sleep quality in students to prevent the development of cardiovascular diseases and early death. Therefore, the purpose of this study was to investigate the relationship between body compositions and hypertension and sleep quality of male students living at the dormitory of Allameh Tabataba'i University.

METHOD

Subjects

This was a descriptive correlational study conducted during August 2018 in a Hemmat dormitory out of all dormitories belong to Allamah Tabatabai was selected randomly. The sample size 168 with a 95% confidence interval and 5% error was determined by the Cochran formula.

A total of 190 out of 300 subjects from all grades and disciplines randomly participated in this study. Then the research aims were presented precisely and all subjects signed the consent form. Also, the researcher assured that the research would be maintained confidential. The subjects' characteristics were collected and recorded by a demographic questionnaire.

The inclusion criteria included: students aged more than 18 years who had given their consent for participation, and experience of living at least 3 years. Excluded if they reported any food consumption for at least 2 hours before sampling, drinking alcohol or a large amount of water, consumption of caffeine-containing substances such as coffee, doing vigorous exercise on sampling day, and aged more than 40-year-old (because of data normalization).

Six subjects were excluded from the study due to the incomplete sleep quality questionnaire and 7 subjects due to the BMI lower than 15kg/m² and measurement error by body composition analysis device 4 subjects due to the constant caffeine consumption and 3 other subjects due to the doing vigorous exercise on the day of sampling. Figure 1 indicates the flow charts of the sampling procedure.




Figure 1

Anthropometric, body composition measurement

The height and weight of subjects were measured and recorded by wall-sticker tape meter and the OMRON digital scanner with 0.01 g sensitivity respectively, with minimal clothing and no shoes. Waist circumference was measured with non-stretching tape at mid-way between the anterior superior iliac crest and the lowest rib. Hip circumference was measured from the side at the greatest point of the buttocks to the nearest 0.1 cm. Waist to hip ratio (WHR) was determined by the corresponding value of waist and hip circumferences and classified into low and high-risk groups according to cutoff point $WHR > 0.9$ in men. Then, body composition indices of subjects were measured and recorded by OMRON (BF511, china) body composition analysis device in a 3-hour fasting state with minimal clothing. Body mass index (BMI) was classified according to the World Health Organization as $< 18.5 \text{ kg/m}^2$ is underweight, $18.5\text{--}24.9 \text{ kg/m}^2$ is normal, $25\text{--}29.9 \text{ kg/m}^2$ is overweight, and $> 30 \text{ kg/m}^2$ is obese. Classification of body fat percentage (BFP) for male aged 18-39 was divided based on manual operation of OMRON Healthcare with ranges of $< 8\%$ low, $8\text{--}19.9\%$ normal, $20\text{--}24.9\%$ high and $> 25\%$ very high ("BF511 body composition monitor," 2015). Also, skeletal muscle mass percentage (SMMP) was classified as $< 33.3\%$ low, $33.3\text{--}39.3\%$

normal, 39.4-44% high and >44.1% very high ("BF511 body composition monitor," 2015).

Sleep quality assessment

To evaluate the students' sleep quality, the Pittsburgh 18 items questionnaire (PSQI) was used. The scoring of this questionnaire is in the form of 3-point Likert scale. The total index of sleep quality measuring by sum of the seven sub-scales. Gaining a score of over 5 in the entire questionnaire means poor sleep quality (Buysse, Reynolds III, Monk, Berman, & Kupfer, 1989).

Blood pressure assessment

Subjects' blood pressure was measured after 10 minutes of rest at the upper left arm in seated position by the OMRON (M2, Vietnam) pressure gauge with an accuracy of 3 mmHg. Hypotension was considered as systolic below 90 mmHg and diastolic below 60 mmHg. Normal blood pressure was defined as 90-119 mmHg (SBP) and 60-79 mmHg (DBP). Subjects with SBP of 120-139 mmHg and DBP of 80-89 mmHg were classified as prehypertensive. Stage-I hypertension was considered as SBP between 140-159 mmHg and DBP between 90-99 mmHg, whereas systolic blood pressure of 160-179 mmHg and diastolic of 100-109 mmHg were classified as stage II hypertension. Hypertensive crisis was determined systolic and diastolic blood over 180 mmHg and 110 mmHg respectively.

Statistical analysis

Data were checked for normality before proceeding to statistics analysis. Descriptive statistics including mean \pm standard deviation, frequencies and percentages were used. Pearson correlation coefficient was used at the significance level of ($P < 0.05$) to find a correlation between anthropometric measures with sleep quality and blood pressure. Data were analyzed using the SPSS software 21 version.

RESULTS

A total of 170 individuals participated in this study, 60 (36%) of whom were overweight. The mean \pm SD age of students was 25.94 ± 3.6 years. Also, 95 (55.88%) of subjects had high or very high body fat. Detailed characteristics of the students are presented in Table 1.

Table 1: demographic, anthropometric and blood pressure measurements

| Variables | Status | Mean±SD | N (%) |
|--|-----------------------------|--------------|--------------|
| Age (years) | | 25.94±3.60 | |
| Height (cm) | | 177.55±6.51 | |
| Weight (kg) | | 75.46±11.38 | |
| BMI (kg/m ²) | | 23.94±3.26 | |
| | Normal | | 104 (61.19%) |
| | Overweight | | 60 (35.29%) |
| | Obese | | 6 (3.52%) |
| Fat percentage (%) | | 20.45±6.73 | |
| | Low | | 6 (3.53%) |
| | Normal | | 69 (40.59%) |
| | High | | 45 (26.47%) |
| | Very high | | 50 (29.41%) |
| Muscle mass (%) | | 39.21±3.91 | |
| | Low | | 5 (2.94%) |
| | Normal | | 89 (52.35%) |
| | High | | 53 (31.18%) |
| | Very High | | 23 (13.53%) |
| WHR | | 0.84±0.06 | |
| | Low risk | | 151 (88.82%) |
| | High risk | | 19 (11.18%) |
| SBP (mmHg) | | 119.97±11.35 | |
| | Normal blood Pressure | | 85 (50%) |
| | Prehypertension | | 78 (45.88%) |
| | High blood pressure stage1 | | 7 (4.12%) |
| DBP (mmHg) | | 75.94±8.89 | |
| | Hypotension | | 5 (2.94%) |
| | Normal blood pressure | | 109 (64.12%) |
| | Prehypertension | | 45 (26.47%) |
| | High blood pressure stage 1 | | 11 (6.47%) |
| BMI: Body mass index, CV: cardiovascular, BP: Blood Pressure | | | |

The mean \pm SD systolic and diastolic blood pressure of subjects was 119.97 ± 11.35 mmHg and 75.94 ± 8.89 mmHg, respectively; however, a total of 78 (45.88%) of the subjects were found to be systolic prehypertensive, and 45 (26.47%) of total participants were found to have diastolic prehypertension.

The results of Pearson correlation coefficient indicated a significant positive correlation between BFP and BMI with both systolic and diastolic blood pressures. In addition, the increase in diastolic blood pressure was associated with an increase in WHR. On the other side, there was a significant and negative correlation between muscle mass and both systolic and diastolic pressure in the students. Also, body composition indices did not show any significant correlation with student's sleep quality (Table 2).

Table 1: The correlation between anthropometric and body composition indices with blood pressure and sleep quality

| Variables | | Fat percentage | Muscle mass | BMI | WHR |
|---------------|---|----------------|-------------|----------|--------|
| Systolic BP | P | +0.000** | -0.001** | +0.000** | 0.147 |
| | r | 0.281 | 0.252 | 0.297 | 0.112 |
| Diastolic BP | P | +0.000** | -0.000** | +0.000** | 0.011* |
| | r | 0.357 | 0.356 | 0.322 | 0.195 |
| Sleep quality | P | 0.764 | 0.846 | 0.933 | 0.906 |
| | r | -0.023 | 0.015 | 0.006 | 0.009 |

* indicates p-value <0.05

** indicating p-value <0.01

DISCUSSION

Since blood pressure is one of the most important risk factors for the metabolic and cardiovascular diseases, the identification and control of aesssss ssssssss nrg ddderrrrr rrrrrrr rrr eeeee can ee eeeeeee as an essential part of health programs. One of the simple and valid methods to tackle this problem is body composition and anthropometric indices assessment. Despite significant progress in health and medicine in recent years, less research has evaluated the relationship between body composition with high blood pressure and sleep quality. The lack of research in this area can endanger the students' health as an important stratum in the community.

In the present study, there was a positive and significant correlation between fat percentage and blood pressure in dormitory students. On the

other hand, in comparison to systolic blood pressure, the association between body fat percentage and diastolic blood pressure was higher, which was consistent with past findings (Moser et al., 2013; Mushengezi & Chillo, 2014). It has been well established that high systolic blood pressure has a significant effect on the incident and development of cardiovascular and metabolic diseases. It has also been reported that an increase in systolic blood pressure leads to an increased risk of intracerebral hemorrhage (stroke caused by bleeding within brain tissue), subarachnoid hemorrhage (the deadliest form of stroke), and stable angina while high diastolic blood pressure is associated with increased risk of Abdominal aortic aneurysm (Rapsomaniki et al., 2014). In addition, it has been reported that high diastolic blood pressure also increases the risk of high systolic blood pressure (Rapsomaniki et al., 2014), so according to our findings (the potent association of body fat percentage with diastolic blood pressure), it can be noted that our subjects with high-fat percentages are at risk of many cardiovascular diseases. Also, our findings indicated that 26.47% of students classified in the high-fat percentage group and 29.41% in the very high-fat percentage group. In fact, 55.88% of the students had fat percentage more than the normal range, which is a very high rate and it can be stated that the risk of hypertension, cardiovascular and metabolic diseases is high in the students of Allameh Tabataba'i University. Our findings are in line with (Wang et al., 2015). An increase in body fat percentage leads to a rise in inflammatory adipokines secretion, which consequently leads to disturbances in secretion and function of nitric oxide (it is very important in regulating blood pressure and vascular function) and decreases its serum levels (Kotsis, Stabouli, Papakatsika, Rizos, & Parati, 2010). Reduction of nitric oxide disturbs endothelial function, resulting in increased stiffness and arterial stenosis, which eventually increases the arterial pressure (Chiolo, Cachat, Burnier, Paccaud, & Bovet, 2007).

In the present study, there was a positive and significant correlation between BMI and students' blood pressure. Also, 38.81% of students were overweight and obese (35.29% overweight and 3.52% obese), which make them susceptible to various diseases such as type 2 diabetes and cardiovascular disease. Our findings in this regard are in line with Deng et al. (2012), Jayedi et al. (Deng et al., 2012; Jayedi, Rashidy-Pour, Khorshidi, & Shab-Bidar, 2018). Regarding the strong and positive

relationship between BMI and body fat, it can be expected that BMI may also increase blood pressure through mentioned direct and indirect mechanisms. It has been reported that high BMI and obesity lead to the adipokines release that is associated with a reduction in efficiency and production of nitric oxide (Kotsis et al., 2010). Nitric oxide is important in vascular dilatation and improvement of blood circulation, and its reduction results in impaired endothelial function which leads to high blood pressure (Chiolero et al., 2007; Kotsis et al., 2010)

In the present study, there was a negative and significant correlation between muscle mass and students' blood pressure. Our findings in this regard, is consistent with studies by Butcher et al. and Benjamin et al. (Benjamin et al., 2017; Butcher et al., 2018). Possible mechanisms of muscle mass effect on blood pressure include reduction of inflammatory factors, improvement of antioxidant enzyme activity and renal function (Butcher et al., 2018). It has also been reported that an increase in muscle mass is often associated with a decrease in body fat percentage, leading to an increase in myostatin (muscle growth inhibitor) inhibition, and consequently results in cardiovascular health, glucose tolerance and endothelial function which ultimately improves blood pressure.

There is a possible mechanism linking muscle mass with blood pressure; to explain more, with an increase in muscle mass, Myostatin, an inhibition factor of muscle growth is blocked, and consequently, it enhances cardiovascular health, glucose tolerance and endothelial function which ultimately leads to blood pressure improvement (Bergen et al., 2015; Ruas et al., 2012).

In the present study, there was a significant positive relationship between waist to hip ratio and diastolic blood pressure. Our findings are in line with the study of Abbaszadeh et al. Factors such as impairment in the renin-angiotensin system, increased stiffness of the arteries, and high activity of sympathetic nervous system have been reported as the most important factors in central obesity-related hypertension (Abbaszadeh et al., 2017).

In the present study, there was no significant relationship between the body composition and sleep quality of dormitory students. Our findings are contradictory to Zhou et al. and Jurado-Fasoli et al. (Jurado-Fasoli et al., 2018; Zhou, Lalani, Banda, & Robinson, 2018). The contradiction reason can be attributed to various factors. In the most

studies, the focus had been on the one or two subcategories of sleep quality such as sleep satisfaction or sleep depth (Harvey, Stinson, Whitaker, Moskovitz, & Virk, 2008), thus the difference in outcome can be due to the different classification and assessing every single factor of sleep quality. On the other hand, in most studies, the sample size were more than 500 subjects, and this may be contributing to the different results (Vargas et al., 2014). Also, the subjects' health status may have a significant impact on the results. In the studies on the overweight, high BMI and high-fat percentage subjects, there was a reverse correlation between these factors and sleep quality (Ye et al., 2018; Zhou et al., 2018). eee ffffeences in eee jjjj ec''' age aa y ee arrrrrr eeannn rrr different results. Therefore, regarding the lack of correlation between obesity and the sleep quality of students, ignoring this could endanger iiiii eeeeeeee eættdd dddddd(Jurado-Fasoli et al., 2018).

CONCLUSIONS

Regarding the wide range of effective factors on sleep quality of young people, indices such as stress, physical activity level, nutritional status and tobacco consumption would be more important indicators than the body composition among dormitory students. The positive role of physical activity to control blood pressure has been proven in many studies, in which most of these studies have reported moderate-intensity aerobic training is the best exercise to modify and control blood pressure and body composition (Ahmed, Blaha, Nasir, Rivera, & Blumenthal, 2012; Boutcher & Boutcher, 2017; Tartibian, Kushkestani, & Nosrani, 2019). Furthermore, improving the diet such as reduction of sodium intake, alcohol consumption, saturated fat intake, and the increase of potassium, and the more consumption of natural and organic substances can be considered as an effective strategy to improve the body composition and prevent the onset of high blood pressure (Bazzano, Green, Harrison, & Reynolds, 2013; Mohanlal, Parsa, & Weir, 2012).

Limitation

As this was a cross-sectional study, it could not clarify the cause-effect relationship between body composition with blood pressure and sleep quality and also sample size was small. Blood pressure was measured only once. It would be more accurate to measure the blood pressure three

times within an interval of at least five minutes and then used its average in the analysis.

Finally, according to the direct relationship between fat percentage and hypertension among students living in the dormitory, it can be concluded that the use of exercise in leisure time and recreational programs to improve body composition can play an important role in the prevention and treatment of hypertension in their future.

Declaration of interest

The authors declare that there is no conflict of interest.

Acknowledgment

We would like to thank all students who participated in the study.

REFERENCES

- Abbaszadeh, F., Sarafraz, N., Atrian, M. K., Sadat, Z., Bagheri, A., & Moravveji, A. (2017). Anthropometric Indices in the Prediction of Hypertension in Female Adolescents. *Iranian Red Crescent Medical Journal*, *19*(11). doi:10.5812/ircmj.14591
- Ahmed, H. M., Blaha, M. J., Nasir, K., Rivera, J. J., & Blumenthal, R. S. (2012). Effects of physical activity on cardiovascular disease. *The American journal of cardiology*, *109*(2), 288-295. doi:10.1016/j.amjcard.2011.08.042
- Bazzano, L. A., Green, T., Harrison, T. N., & Reynolds, K. (2013). Dietary approaches to prevent hypertension. *Current hypertension reports*, *15*(6), 694-702. doi:10.1007/s11906-013-0390-z
- Benjamin, E. J., Blaha, M. J., Chiuve, S. E., Cushman, M., Das, S. R., Deo, R., ... & Jiménez, M. C. (2017). Heart disease and stroke statistics-2017 update: a report from the American Heart Association. *circulation*, *135*(10), e146-e603. doi:10.1161/cir.0000000000000485
- Bergen, H. R., Farr, J. N., Vanderboom, P. M., Atkinson, E. J., White, T. A., Singh, R. J., ... & LeBrasseur, N. K. (2015). Myostatin as a mediator of sarcopenia versus homeostatic regulator of muscle mass: insights using a new mass spectrometry-based assay. *Skeletal muscle*, *5*(1), 21. doi:10.1186/s13395-015-0047-5
- Bixler, E. (2009). Sleep and society: an epidemiological perspective. *Sleep medicine*, *10*, S3-S6. doi:10.1016/j.sleep.2009.07.005

- BF511, O. (2015). Body Composition Monitor. *Instruction Manual, All for Healthcare*, Retrieved from Omron Healthcare <https://gzhls.at/blob/ldb/e/6/f/c/be4cc09431067d512bd0a318aef6884c3008.pdf>.
- Bloch, M. J. (2016). Worldwide prevalence of hypertension exceeds 1.3 billion. *Journal of the American Society of Hypertension: JASH*, 10(10), 753. doi:10.1016/j.jash.2016.08.006
- Boutcher, Y. N., & Boutcher, S. H. (2017). Exercise intensity and hypertension: wttt '> *Journal of human hypertension*, 31(3), 157. doi:10.1038/jhh.2016.62
- Buboltz Jr, W., Jenkins, S. M., Soper, B., Woller, K., Johnson, P., & Faes, T. (2009). Sleep habits and patterns of college students: an expanded study. *Journal of College Counseling*, 12(2), 113-124. doi:10.1002/j.2161-1882.2009.tb00109.x
- Butcher, J. T., Mintz, J. D., Larion, S., Qiu, S., Ruan, L., Fulton, D. J., & Stepp, D. W. (2018). Increased muscle mass protects against hypertension and renal injury in obesity. *Journal of the American Heart Association*, 7(16), e009358. doi:10.1161/JAHA.118.009358
- Buysse, D. J., Reynolds III, C. F., Monk, T. H., Berman, S. R., & Kupfer, D. J. (1989). The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry research*, 28(2), 193-213. doi:10.1016/0165-1781(89)90047-4
- Chiolero, A., Cachat, F., Burnier, M., Paccaud, F., & Bovet, P. (2007). Prevalence of hypertension in schoolchildren based on repeated measurements and association with overweight. *Journal of hypertension*, 25(11), 2209-2217. doi:10.1097/HJH.0b013e3282ef48b2
- laary, ,, Wll ter, G., & aakk,,, , . (2011). oot a.wyys mntt h aailigg:: Mental health issues associated with the transition from high school to college. *Issues in mental health nursing*, 32(4), 250-254. doi:10.3109/01612840.2010.548906
- Deng, W. W., Wang, J., Liu, M. M., Wang, D., Zhao, Y., Liu, Y. Q., ... & Dong, G. H. (2012). Body mass index compared with abdominal obesity indicators in relation to prehypertension and hypertension in adults: the CHPSNE study. *American journal of hypertension*, 26(1), 58-67. doi:10.1093/ajh/hps001
- Ettehad, D., Emdin, C. A., Kiran, A., Anderson, S. G., Callender, T., Emberson, J., ... & Rahimi, K. (2016). Blood pressure lowering for prevention of cardiovascular disease and death: a systematic review and meta-analysis. *The Lancet*, 387(10022), 957-967. doi:10.1016/S0140-6736(15)01225-8

- Harvey, A. G., Stinson, K., Whitaker, K. L., Moskovitz, D., & Virk, H. (2008). The subjective meaning of sleep quality: a comparison of individuals with and without insomnia. *Sleep, 31*(3), 383-393. doi:10.1093/sleep/31.3.383
- Hypertension. (2019, September 13). Retrieved January 5, 2020, from <https://www.who.int/news-room/fact-sheets/detail/hypertension>.
- Jayedi, A., Rashidy-Pour, A., Khorshidi, M., & Shab-Bidar, S. (2018). Body mass index, abdominal adiposity, weight gain and risk of developing hypertension: a systematic review and dose-response meta-analysis of more than 2.3 million participants. *Obesity reviews, 19*(5), 654-667. doi:10.1111/obr.12656
- Jurado-Fasoli, L., Amaro-Gahete, F., De-la-O, A., Dote-Montero, M., Gutiérrez, Á., & Castillo, M. (2018). Association between Sleep Quality and Body Composition in Sedentary Middle-Aged Adults. *Medicina, 54*(5), 91. doi:10.3390/medicina54050091
- Kahlhöfer, J., Karschin, J., Breusing, N., & Bosy-Westphal, A. (2016). Relationship between actigraphy-assessed sleep quality and fat mass in college students. *Obesity, 24*(2), 335-341. doi:10.1002/oby.21326
- Kishi, S., Teixido-Tura, G., Ning, H., Venkatesh, B. A., Wu, C., Almeida, A., ... & Gidding, S. S. (2015). Cumulative blood pressure in early adulthood and cardiac dysfunction in middle age: the CARDIA study. *Journal of the American College of Cardiology, 65*(25), 2679-2687. doi:10.1016/j.jacc.2015.04.042
- Kotsis, V., Stabouli, S., Papakatsika, S., Rizos, Z., & Parati, G. (2010). Mechanisms of obesity-induced hypertension. *Hypertension research, 33*(5), 386. doi:10.1038/hr.2010.9
- Kronholm, E., Partonen, T., Laatikainen, T., Peltonen, M., Härmä, M., Hublin, C., ... & Valve, R. (2008). Trends in self-reported sleep duration and insomnia-related symptoms in Finland from 1972 to 2005: a comparative review and re-analysis of Finnish population samples. *Journal of sleep research, 17*(1), 54-62. doi:10.1111/j.1365-2869.2008.00627.x
- Mohanlal, V., Parsa, A., & Weir, M. R. (2012). Role of dietary therapies in the prevention and treatment of hypertension. *Nature Reviews Nephrology, 8*(7), 413. doi:10.1038/nrneph.2012.91
- Moser, D. C., Giuliano, I. D. C. B., Titski, A. C. K., Gaya, A. R., Coelho-e-Silva, M. J., & Leite, N. (2013). Anthropometric measures and blood pressure in school children. *Jornal de pediatria, 89*(3), 243-249. doi:10.1016/j.jpmed.2012.11.006

- Mushengezi, B., & Chillo, P. (2014). Association between body fat composition and blood pressure level among secondary school adolescents in Dar es Salaam, Tanzania. *The Pan African medical journal*, *19*. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4405073/>
- Nyombi, K. V., Kizito, S., Mukunya, D., Nabukalu, A., Bukama, M., Lunyera, J., ... & Kalyesubula, R. (2016). High prevalence of hypertension and cardiovascular disease risk factors among medical students at Makerere University College of Health Sciences, Kampala, Uganda. *BMC research notes*, *9*(1), 110. Retrieved from <https://bmresnotes.biomedcentral.com/articles/10.1186/s13104-016-1924-7>
- Oori, M. J., Mohammadi, F., Norozi, K., Fallahi-Khoshknab, M., Ebadi, A., & Gheshlagh, R. G. (2019). Prevalence of HTN in Iran: Meta-analysis of Published Studies in 2004-2018. *Current hypertension reviews*, *15*(2), 113-122. doi:10.2174/1573402115666190118142818
- Patel, H. C., Hayward, C., Ozdemir, B. A., Rosen, S. D., Krum, H., Lyon, A. R., ... & Di Mario, C. (2015). Magnitude of blood pressure reduction in the placebo arms of modern hypertension trials: implications for trials of renal denervation. *Hypertension*, *65*(2), 401-406. doi:10.1161/HYPERTENSIONAHA.114.04640
- Rahe, C., Czira, M. E., Teismann, H., & Berger, K. (2015). Associations between poor sleep quality and different measures of obesity. *Sleep medicine*, *16*(10), 1225-1228. doi:10.1016/j.sleep.2015.05.023
- Rapsomaniki, E., Timmis, A., George, J., Pujades-Rodriguez, M., Shah, A. D., Denaxas, S., ... & Williams, B. (2014). Blood pressure and incidence of twelve cardiovascular diseases: lifetime risks, healthy life-years lost, and age-specific associations in 1·25 million people. *The Lancet*, *383*(9932), 1899-1911. doi:10.1016/S0140-6736(14)60685-1
- Ruas, J. L., White, J. P., Rao, R. R., Kleiner, S., Brannan, K. T., Harrison, B. C., ... & Lanza, I. R. (2012). A PGC-1 α pathway regulates skeletal muscle hypertrophy. *Cell*, *151*(6), 1319-1331. doi:10.1016/j.cell.2012.10.050
- Sin, C. W., Ho, J. S., & Chung, J. W. (2009). Systematic review on the effectiveness of caffeine abstinence on the quality of sleep. *Journal of Clinical Nursing*, *18*(1), 13-21. doi:10.1111/j.1365-2702.2008.02375.x
- Suen, L. K., Ellis Hon, K. L., & Tam, W. W. (2008). Association between sleep behavior and sleep-related factors among university students in Hong Kong. *Chronobiology international*, *25*(5), 760-775. doi:10.1080/07420520802397186

- Tartibian, B., Kushkestani, M., & Ebrahimpour Nosrani, S. (2019). The Effect of 12-Week Endurance Training on Lipid Profiles and Fat Percentage of Overweight Girls. *New Approaches in Sport Sciences*, 1(1), 189-200. doi:10.22054/NASS.2019.10134
- Vargas, P. A., Flores, M., & Robles, E. (2014). Sleep quality and body mass index in college students: the role of sleep disturbances. *Journal of American College Health*, 62(8), 534-541. doi:10.1080/07448481.2014.933344
- Wang, Z., Zeng, X., Chen, Z., Wang, X., Zhang, L., Zhu, M., & Yi, D. (2015). Association of visceral and total body fat with hypertension and prehypertension in a middle-aged Chinese population. *Journal of hypertension*, 33(8), 1555-1562. 10.1097/HJH.0000000000000602
- Wells, S., Riddell, T., Kerr, A., Pylypchuk, R., Chelimo, C., Marshall, R., ... & Grey, C. (2015). Cohort profile: the PREDICT cardiovascular disease cohort in New Zealand primary care (PREDICT-CVD 19). *International journal of epidemiology*, 46(1), 22-22. doi:10.1093/ije/dyv312
- Ye, S., Zhu, C., Wei, C., Yang, M., Zheng, W., Gan, D., & Zhu, S. (2018). Associations of Body Composition with Blood Pressure and Hypertension. *Obesity*, 26(10), 1644-1650. doi:10.1002/oby.22291
- Zhou, M., Lalani, C., Banda, J. A., & Robinson, T. N. (2018). Sleep duration, timing, variability and measures of adiposity among 8-to 12-year-old children with obesity. *Obesity science & practice*, 4(6), 535-544. doi:10.1002/osp4.303