



**Research Paper: Associations between Physical Activity with Self-Esteem and Perceived Motor Competence among Children with Developmental Coordination Disorder**



**Tayebeh Baniasadi<sup>1</sup>, Sedigheh Khajaeiflaton Mofrad<sup>2</sup>, Forough ShafaeianFard<sup>3</sup>**

<sup>1</sup> Visiting Scholar, Indiana University, School of Public Health, Department of Kinesiology, USA

<sup>2</sup> Department of Physical Education, Farhangian University, Gorgan, Iran.

<sup>3</sup> M.A Student, Iran

**Citation:** Baniasadi, T., Khajaeiflaton Mofrad, S., ShafaeianFard, F. (2022). Associations between Physical Activity with Self-Esteem and Perceived Motor Competence among Children with Developmental Coordination Disorder. *Journal of Modern Psychology*, 2(2),1-10. <https://doi.org/10.22034/JMP.2022.357946.1038>

 <https://doi.org/10.22034/JMP.2022.357946.1038>

**Article info:**

**Received date:**

15 Jan. 2022

**Accepted date:**

22 Mar. 2022

**Keywords:**

DCD, Perceived motor competence, Physical activity, Self-esteem

**Abstract**

Evidence shows that physical activity (PA) is associated with better self-esteem and perceived motor competence in typically developing children. Nonetheless, associations between PA with self-esteem and perceived motor competence among children with developmental coordination disorder (DCD) has received very little attention in the literature. As such, the purpose of this study was to survey the associations between PA with self-esteem and perceived motor competence among children with DCD. A correlational approach was used in this study. Participants were forty-nine children with DCD (mean age of 8.85 years old) from special schools. Physical Activity Questionnaire for Older Children was used for assessing PA. Self-Perception Profile for Children was used to assess self-esteem and perceived motor competence. To analyze data, we used Pearson correlation test and regression analysis. Descriptive results showed that our sample participate in very low amount of PA. In addition, they have low levels of self-esteem and perceived motor competence. Furthermore, PA was significantly and directly associated with both self-esteem and perceived motor competence. Finally, PA has significantly and directly predicted both self-esteem and perceived motor competence. Our findings emphasize on benefits of PA for improving psychological status of children with DCD. Thus, it seems necessary to find out proper strategies and interventions for increase the level of PA in this population.

**\* Corresponding author:**

Tayebeh Baniasadi

**Address:** Indiana University, School of Public Health, Department of Kinesiology, USA

**Tel:** +98 (911) 222 3333

**E-mail:** [tbaniasa@iu.edu](mailto:tbaniasa@iu.edu)



© 2022, The Author(s). Published by *Rahman Institute of Higher Education*. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)

## 1. Introduction

Developmental coordination disorder (DCD), also known as developmental motor coordination disorder, developmental dyspraxia or simply dyspraxia, is a lifelong condition that affects learning and executing motor skills and coordination. DCD is not a learning disorder, but it can impact learning. Children with DCD struggle with physical tasks and activities they need to do both in and out of school (Barnett et al. 2019; Cantell et al. 2003; Fogel et al. 2021; Gomez et al. 2017; Tseng et al. 2007). Several studies have demonstrated that children with DCD show poor motor competence in everyday self-care and academic activities such as catching, using scissors, handwriting, riding a bike and participating in sports (Barnett et al. 2019; Cantell et al. 2003; Fogel et al. 2021; Gomez et al. 2017; O'Dea & Connell, 2016; Saban et al. 2014; Tseng et al. 2007). Some studies have also shown that for most individuals with DCD difficulties can persist into adulthood and affect the learning and performance of new motor skills such as driving (Barnhart et al. 2003; Cantell et al. 1994, 2003). Moreover, other line of research has demonstrated that individuals with DCD show low levels of self-esteem, self-confidence, and mental health (Cairney et al. 2005). However, some factors might can positively influence poor physical and psychological components associated with individuals with DCD. An influential factor might be engaging in regular physical activity (PA).

PA refers to any voluntary bodily movement executed by skeletal muscles that requires energy expenditure. Some of physical activities included homework, gardening, sport, walking, etc. (Caspersen

et al. 1985; Thivel et al. 2018; Ghorbani et al. 2021). PA leads to numerous physical and mental health benefits in all age categories, including children. Some of these benefits encompass improvement in brain health, helping in weight management, reduction of the risk of disease, strengthening bones and muscles, improving quality of life, and improving ability to do everyday activities (Abdoshahi, Gholami, Naeimikia, 2022; Basterfield et al. 2021; Dana & Christodoulides, 2019; Dana et al. 2021; Hashemi Motlagh, BaniAsadi, Chaharbaghi, & Moradi, 2022; Gholami & Rostami, 2021; Ghorbani et al. 2020, 2021; Lahart et al. 2019; Mohammad Gholinejad, Hojjati, & Ghorbani, 2019; Mohammadi, Nafei, Baniasadi, & Chaharbaghi, 2022; Naeimikia, Izanloo, Gholami, & Ahar, 2018; Naeimikia & Gholami, 2018, 2020; Schwartz et al. 2019; Tremblay et al. 2011; Wafa et al. 2016; Yaali, Naeimi Kia, Gholami, 2018; Zhang et al. 2021). Due to so many benefits of PA, world health organization (WHO) recommends a proper amount of PA for children aged 7 to 18 years old (i.e., at least 60 minutes of moderate-to-vigorous PA across the week) (Bull et al. 2020). Regarding children with DCD, however, it has been shown that they do not follow WHO guidelines (Cermak et al. 2015; Steenbergen et al. 2020). As well, children with DCD demonstrated significantly reduced PA, increased sedentary behavior, poorer fitness and increased overweight compared with typically developing children (Cermak et al. 2015; Steenbergen et al. 2020), which may be due to their physical limitations. It should be noted that lower amount of PA in individuals with DCD can potentially lead to various negative consequences such as enhancing the risk of chronic diseases such

as type 2 diabetes and cardiovascular disease (Kinne et al. 2004; Rimmer et al. 2007).

Furthermore, in healthy children, research has shown that children who participate in regular PA have higher levels of self-esteem and perceived motor competencies (Cantell et al. 2008; Haga, 2009). As mentioned earlier, being DCD results in low self-esteem and other psychological components (Cairney et al. 2005). However, associations between PA and psychological variables have not been investigated in children with DCD. Therefore, the aim of this study was to investigate the associations between PA with self-esteem and perceived motor competence among children with DCD. It was hypothesized that children who PA will positively affect self-esteem and perceived motor competence in children with DCD.

## 2. Methods

### 2.1 Participants

This study was conducted based on correlational research method. The sample included 49 children with DCD between 8 to 10 years old (mean 8.85 years old) who attended in special schools. All participants have voluntarily attended in the study. The parents of children gave informed consents. Protocol of this study was in accordance with ethical guidelines of declaration of Helsinki. According to school's office, all children were already diagnosed as DCD. In this study, an experienced examiner assessed the symptoms of DCD in children using the American Psychiatric Association's (2000) *Diagnostic and Statistical Manual of Mental Disorders Text Revision. 4th ed*, too.

## 2.2 Measures

**2.2.1 Physical Activity:** Physical Activity Questionnaire for Older Children (PAQ-C) was used for measuring PA of children with DCD. The PAQ-C is a self-administered, 7-day recall instrument. It assesses general levels of PA throughout the elementary school year for students approximately 8 to 14 years of age. The PAQ-C contains nine items, each scored on a 5-point scale (Crocker et al. 1997). Reliability of PAQ-C was measured in this study where Cronbach's alpha coefficient was 0.85.

**2.2.2 Self-Esteem and Perceived Motor Competence:** In this study, the Self-Perception Profile for Children (Harter, 1985) was used to assess self-esteem and perceived motor competence. This is a commonly used scale with good validity and reliability for use with a child population. It comprises a 36-item self-completed questionnaire measuring perceived competence in five domains (scholastic competence, social acceptance, athletic competence, physical appearance and behavioral conduct) and feelings of global self-worth. Each domain consists of six paired statements. Respondents are asked to select the statement that best describes them and then to say whether it is 'really true' or 'sort of true' for them. In this study, for evaluating self-esteem, we used self-worth was used. As such, for assessing perceived motor competence, we used the part of athletic competence. In this study, reliability of these scales was measured where Cronbach's alpha coefficients were 0.93 and 0.90 for self-esteem and perceived motor competence, respectively.

### 2.3 Data analysis

To describe research variables, we used mean and standard deviation. Kolmogorov-Smirnov test was used for measuring the normality of data. Pearson correlation test was used to measure the associations between research variables. Finally, regression analysis was used to investigate whether PA predicts self-esteem and perceived motor competence in children with DCD. SPSS software version 26 was used to analyze the data. P-value was set at  $P < 0.05$ .

### 3. Results

#### 3.1 Descriptive Results

Mean and standard deviations and relationships between of research variables are shown in Table 1. Descriptive results showed that the level of PA was very low in children with DCD. In addition, children with ASD had low scores in self-esteem and perceived motor competence. Results of Kolmogorov-Smirnov tests showed that our data were normally distributed (all  $P > 0.05$ ).

Table 1

*Mean, standard deviation and relation between research variables*

Variables	M	SD	1	2	3
1. Physical Activity	1.01	0.53	-		
2. Self-Esteem	6.82	4.19	0.69***	-	
3. Perceived Motor Competence	7.48	5.22	0.51***	0.74***	-

Results in Table 1 demonstrated that there were significant associations between PA and self-esteem among children with DCD ( $p=0.000$ ). In addition, PA was significantly associated with perceived motor competence among children with DCD ( $p=0.000$ ).

#### 3.2 Results of Regression Analysis

Results of regression analysis are presented in Table 2. As observed, PA has directly predicted self-esteem among children with DCD ( $p=0.000$ ). Also, PA has directly predicted perceived motor competence among children with DCD ( $p=0.000$ ).

Table 2

The results of multiple regression analysis for predicting self-esteem and perceived motor competence by PA

criteria variable	B	SE	Beta	T	Sig	Tolerance	VIF
Self-esteem			0.582	4.697			
Perceived motor competence			0.493	3.415			

$R=0.508$   $R^2=0.285$   $F=8.694$   $P \leq 0.001$

$R=0.465$   $R^2=0.216$   $F=6.128$   $P \leq 0.001$

### 4. Discussion

Previous studies have shown that PA has direct relationship with psychological

status (e.g., self-esteem and perceived motor competence) of typically developing children (Cantell et al. 2008; Haga, 2009). However, associations between PA and



psychological status of children in special groups such as DCD have been not surveyed. Thus, the purpose of this study was to investigate the associations between PA with self-esteem and perceived motor competence among children with DCD. It was hypothesized that PA has significant associations with self-esteem and perceived motor competence among children with DCD.

First of all, results of descriptive data showed that the children in this study had very low level of PA, which are in line with the findings of previous studies (Cermak et al. 2015; Steenbergen et al. 2020), indicating that children with DCD engage in very low amount of PA, which may be because of their physical limitations. Furthermore, the level of self-esteem and perceived motor competence were very low in children with DCD in this study, which are consistent with the findings of previous studies (Barnett et al. 2019; Cantell et al. 2003; Fogel et al. 2021; Gomez et al. 2017; O'Dea & Connell, 2016; Saban et al. 2014; Tseng et al. 2007), indicating psychological problems among this population. Psychological problems in individuals with DCD might be a consequence of their physical limitation and barriers. They do not engage in various physical activities which make them distancing from their peers. Most of children with DCD experience loneliness and it is possible that their mental health is negatively affected. According to the results of this study, it seems necessary to adopt appropriate strategies and interventions to enhance the level of PA and mental health among this population.

Additionally, the results of this study revealed that PA was significantly

associated with both self-esteem and perceived motor competence among children with DCD. Also, the results of regression analysis showed that higher levels of PA may directly predict higher levels of both self-esteem and perceived motor competence in children with DCD. These results confirm the findings of previous studies on typically developing children (Cantell et al. 2008; Haga, 2009), indicating the positive role played by PA in improving psychological status among children with DCD. As mentioned earlier, children with DCD suffer from poor motor competence. Our findings indicate that if they participate in more PA, it can be resulted in enhancing the perception of their abilities in performing motor skills. In addition, participating in regular PA can be resulted in increasing the level of their self-esteem, which is an important psychological variable in mental health. Thus, Therefore, it can be proposed that children with DCD who participate in regular PA have better mental health compared with those who do not participate in regular PA. The findings of present study also generalize the positive benefits of PA among children with DCD (Abdoshahi, Gholami, Naeimikia, 2022; Basterfield et al. 2021; Dana & Christodoulides, 2019; Dana et al. 2021; Hashemi Motlagh, BaniAsadi, Chaharbaghi, & Moradi, 2022; Gholami & Rostami, 2021; Ghorbani et al. 2020, 2021; Lahart et al. 2019). In early stages of childhood, participating in regular PA provides optimal condition for enhancing motor proficiency in children, especially those in special groups such as DCD (BaniAsadi et al. 2019; Chaharbaghi et al. 2022; Mohammad Gholinejad, Hojjati, & Ghorbani, 2019; Mohammadi, Nafei, BaniAsadi, & Chaharbaghi, 2022;

Naeimikia, Izanloo, Gholami, & Ahar, 2018; Naeimikia & Gholami, 2018, 2020; Schwartz et al. 2019; Tremblay et al. 2011; Wafa et al. 2016; Yaali, Naeimi Kia, Gholami, 2018; Zhang et al. 2021). As such, it can be proposed that enhancing the level of PA in children with DCD results in better psychological status.

As a limitation to this study, it can be stated that using questionnaire for measuring PA has self-reporting bias (Ghorbani et al. 2021). Thus, future studies should use modern devices for measuring PA for collecting precise data. Also, we have included only 49 children with DCD in the study, however, it seems a relatively small size. Future studies should use larger sample size for collecting more reliable data.

## 5. Conclusion

To summarize, the present study adds some new findings into the literature by showing that children who have higher amount of PA have better mental status (e.g., self-esteem and perceived motor competence). Along with the fact that our sample had very low levels of PA and psychological status, it seems that participating in regular PA can act as a proper strategy for diminish the conditions of DCD among this population. Thus, it is essential to find out the ways to increase the level of PA among children with DCD.

## Acknowledgement

The authors are thankful to all the people who participated in this study and contributed to facilitate the research process.

## Conflict of interest

The Authors declare that there is no conflict of interest with any organization. Also, this research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## References

- Abdoshahi, M., Gholami, A., & Naeimikia, M. (2022). The correlation of autonomy support with intrinsic motivation, anxiety, and intention to do physical activities in children. *International Journal of Pediatrics*, 10(3), 15623-15629. <https://dx.doi.org/10.22038/ijp.2022.63021.4810>
- American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders, Text Revision*. 4th ed. Washington, DC: American Psychiatric Association. <https://www.amazon.com/Diagnostic-Statistical-Disorders-Revision-DSM-IV-TR/dp/0890420254>
- Barnett, A.L., Law, C., & Stuart, N. (2019). *Developmental progression in DCD*. In: Barnett AL, Hill EL, editors. Understanding motor behavior in developmental coordination disorder. London: Routledge; p. 28–51.
- Baniasadi, T., Namazi Zadeh, M., Sheikh, M. (2019). The effects of balance training and focus of attention on sway in postural and supra-postural tasks in the elderly

- population. *Motor Behavior*, 11(36), 89-104. [https://mbj.ssric.ac.ir/article\\_1506.html](https://mbj.ssric.ac.ir/article_1506.html)
- Barnhart, R.C., Davenport, M.J., Epps, S.B., & Nordquist, V.M. (2003). Developmental coordination disorder. *Physical Therapy*, 83, 722-731.
- Basterfield, L., Burn, N.L., Galna, B., Karoblyte, G., & Weston, K.L. (2021). The association between physical fitness, sports club participation and body mass index on health-related quality of life in primary school children from a socioeconomically deprived area of England. *Preventive Medicine Reports*, 24, 101557. <https://doi.org/10.1016/j.pmedr.2021.101557>
- Bull, F.C., et al. (2020). World Health Organization 2020 guidelines on physical activity and sedentary behavior. *British Journal of Sports Medicine*, 54(24), 1451-1462. <http://dx.doi.org/10.1136/bjsports-2020-102955>
- Cairney, J., Hay, J.A., Faight, B.E., Wade, T.J., Corna, L., & Flouris, A. (2005). Developmental coordination disorder, generalized self-efficacy toward physical activity, and participation in organized and free play activities. *The Journal of Pediatrics*, 147(4), 515-520. <https://doi.org/10.1016/j.jpeds.2005.05.013>
- Cantell, M.H., Smyth, M.M., & Ahonen, T.P. (1994). Clumsiness in adolescence: Educational, motor, and social outcomes of motor delay detected at 5 Years. *Adapted Physical Activity Quarterly*, 11(2), 115-129. <https://doi.org/10.1123/apaq.11.2.115>
- Cantell, M., Smyth, M.M., & Ahonen, T.P. (2003). Two distinct pathways for developmental coordination disorder: persistence and resolution. *Human Movement Science*, 22, 413-431. <https://doi.org/10.1016/j.humov.2003.09.002>
- Cantell, M., Crawford, S.G., & Doyle-Baker, P.K. (2008). Physical fitness and health indices in children, adolescents and adults with high or low motor competence. *Human Movement Science*, 27, 344-362. <https://doi.org/10.1016/j.humov.2008.02.007>
- Caspersen, C. J., Powell, K. E., & Christenson, G. M. (1985). Physical activity, exercise, and physical fitness: Definitions and distinctions for health-related research. *Public Health Reports*, 100, 126-131. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1424733/>
- Cermak, S.A., et al. (2015). Participation in physical activity, fitness, and risk for obesity in children with developmental coordination disorder: A cross-cultural study. *Occupational Therapy International*, 22(4), 163-173.
- Chaharbaghi, Z., Hosseini, F., Baniasadi, T., Moradi, L., Dana, A. (2022). Impact of physical activity on resilience among teenage girls during the COVID-19 pandemic: a mediation by self-esteem. *Women's Health Bulletin*, 9(2), 80-85. <https://dx.doi.org/10.30476/whb.2022.94451.1166>
- Crocker, P.R.E., Bailey, D.A., Faulkner, R.A., Kowalski, K.C., & McGrath, R. (1997). Measuring general levels of physical activity: Preliminary evidence for the Physical Activity Questionnaire for Older Children. *Medicine & Science in Sports & Exercise*, 29, 1344-1349. <https://doi.org/10.1097/00005768-199710000-00011>
- Dana, A., & Christodoulides, E. (2019). The effects of a period of selected physical activity on improving manipulative and locomotor skills of children with neuropsychological learning disabilities, *The Journal of Rehabilitation Sciences & Research*, 7, 25-30.

- <https://doi.org/10.30476/jrsr.2019.81592.1006>
- Dana, A., Khajehaflatan, S., Salehian, M., & Sarvari, S. (2021). Effects of an intervention in online physical education classes on motivation, intention, and physical activity of adolescents during the COVID-19 pandemic. *International Journal of School Health*, 8(3), 141-149. <https://dx.doi.org/10.30476/intjsh.2021.91103.1145>
- Fogel, Y., Stuart, N., Joyce, T., & Barnett, A.L. (2021). Relationships between motor skills and executive functions in developmental coordination disorder (DCD): A systematic review. *Scandinavian Journal of Occupational Therapy*, 28, 1-13. <https://doi.org/10.1080/11038128.2021.2019306>
- Gholami, A., & Rostami, S. (2021). Effect of a fun virtual purposeful active play program on children's physical fitness during home quarantine due to the outbreak of Covid-19. *Motor Behavior*, 13(44), 171-190. <https://dx.doi.org/10.22089/mbj.2021.10913.1980>
- Ghorbani S, Afshari M, Eckelt M, Dana A, & Bund A. (2021). Associations between physical activity and mental health in Iranian adolescents during the COVID-19 pandemic: An accelerometer-based study. *Children*, 8(11), 1022. <https://doi.org/10.3390/children8111022>
- Ghorbani, S., Rezaeeshirazi, R., Shakki, M., Noohpishah, S., & Farzanegi, P. (2020). The role of BMI, physical activity and the use of electronic device in the status of trunk abnormalities in male adolescents. *Journal of Gorgan University of Medical Sciences*, 22(3), 129-136. <http://goums.ac.ir/journal/article-1-3676-en.html>
- Gomez, A., et al. (2017). Numerical abilities of school-age children with developmental coordination disorder (DCD): a behavioral and eye-tracking study. *Human Movement Science*, 55, 315-326. <https://doi.org/10.1016/j.humov.2016.08.008>
- Haga, M. (2009). Physical fitness in children with high motor competence is different from that in children with low motor competence. *Physical Therapy*, 89(10), 1089-1097. <https://doi.org/10.2522/ptj.20090052>
- Harter, S. (1985). *Manual for the Self-Perception Profile for Children*, Denver: University Press.
- Hashemi Motlagh, S., BaniAsadi, T., Chaharbaghi, Z., & Moradi, L. (2022). The effects of socioeconomic status on physical activity in children: Mediating role of motivation. *International Journal of Pediatrics*, 10(8), 16538-16544. <https://dx.doi.org/10.22038/ijp.2022.63421.4834>
- Kinne, S., Patrick, D. L., & Doyle, D. L. (2004). Prevalence of secondary conditions among people with disabilities. *American Journal of Public Health*, 94(3), 443-445. <https://ajph.aphapublications.org/doi/full/10.2105/AJPH.94.3.443>
- Lahart I, Darcy P, Gidlow C, & Calogiuri G. (2019). The Effects of Green Exercise on Physical and Mental Wellbeing: A Systematic Review. *International Journal of Environmental Research & Public Health*, 16(8), 1352. <https://doi.org/10.3390/ijerph16081352>
- Mohammad Gholinejad, P., Hojjati, H., & Ghorbani, S. (2019). The effect of aerobic exercise on body composition and muscle strength of female students at elementary schools of Ali Abad Katoul in 2018. *International Journal of School Health*, 6(4), 27-33. <https://dx.doi.org/10.30476/intjsh.2019.45892>
- Mohammadi, H., Nafei, H., BaniAsadi, T., & Chaharbaghi, Z. (2022). Accelerometer-



- based physical activity and health-related quality of life in children with ADHD. *International Journal of Pediatrics*, 10(7), 16362-16369. <https://dx.doi.org/10.22038/ijp.2022.63699.4847>
- Naeimikia, M., Izanloo, Z., Gholami, A., & Ahar, S. (2018). The effect of walking training with cognitive loading on gait indicators related to balance in elderly males. *Journal of Geriatric Nursing*, 4(3), 43-53. <https://jgn.medilam.ac.ir/article-1-255-fa.pdf>
- Naeimikia, M., & Gholami, A. (2018). Effect of walking training on artificial cobblestone mats on gait spatiotemporal parameters for the elderly women. *Motor Behavior*, 9(30), 71-86. <https://dx.doi.org/10.22089/mbj.2018.3683.1447>
- Naeimikia, M., & Gholami, A. (2020). Effect of physical activity on the level of perceived mental pressure during home quarantine due to coronavirus outbreak. *Journal of Rehabilitation Medicine*, 9(3), 217-224. [http://www.medrehab.sbm.ac.ir/article\\_1101156.html?lang=en](http://www.medrehab.sbm.ac.ir/article_1101156.html?lang=en)
- O'Dea, A., & Connell, A. (2016). Performance difficulties, activity limitations and participation restrictions of adolescents with developmental coordination disorder (DCD). *British Journal of Occupational Therapy*, 79, 540-549. <https://doi.org/10.1177/0308022616643100>
- Rimmer, J.H., Rowland, J.L., & Yamaki, K. (2007). Obesity and secondary conditions in adolescents with disabilities: Addressing the needs of an underserved population. *Journal of Adolescent Health*, 41(3), 224-229. <https://doi.org/10.1016/j.jadohealth.2007.05.005>
- Saban, M.T., Ornoy, A., & Parush, S. (2014). Executive function and attention in young adults with and without developmental coordination disorder-a comparative study. *Research in Developmental Disabilities*, 35, 2644-2650. <https://doi.org/10.1016/j.ridd.2014.07.002>
- Schwartz, J., Rhodes, R., Bredin, S., Oh, P., & Warburton, D. (2019). Effectiveness of approaches to increase physical activity behavior to prevent chronic disease in adults: A brief commentary. *Journal of Clinical Medicine*, 8(3), 295. <https://doi.org/10.3390/jcm8030295>
- Steenbergen, B., Bekhuis, H. & van Abswoude, F. (2020). Promoting participation in DCD: Physical activity levels and the social network. *Current Developmental Disorders Reports*, 7, 43-47. <https://doi.org/10.1007/s40474-020-00193-y>
- Thivel, D., Tremblay, A., Genin, P.M., Panahi, S., Rivière, D., & Duclos, M. (2018). Physical activity, inactivity, and sedentary behaviors: Definitions and implications in occupational health. *Frontiers in Public Health*, 6, 288. <https://www.frontiersin.org/articles/10.3389/fpubh.2018.00288/full>
- Tremblay, M.S., et al. (2011). Systematic review of sedentary behavior and health indicators in school-aged children and youth. *International Journal of Behavioral Nutrition & Physical Activity*, 8, 98. <https://doi.org/10.1186/1479-5868-8-98>
- Tseng, M.H., et al. (2007). Cooccurrence of problems in activity level, attention, psychosocial adjustment, reading and writing in children with developmental coordination disorder. *International Journal of Rehabilitation Research*, 30, 327-332. <https://doi.org/10.1097/mrr.0b013e3282f144c7>
- Wafa, S.W., et al. (2016). Association between physical activity and health-related quality of life in children: a cross-sectional study.

- Health & Quality of Life Outcomes*, 4, 71. <https://doi.org/10.1186/s12955-016-0474-y>
- Yaali, R., Naeimi Kia, M., & Gholami, A. (2018). Effect of weight transfer training on static and dynamic balance of older women. *Research in Sport Management & Motor Behavior*, 8(16), 47-59. [https://jrsm.khu.ac.ir/browse.php?a\\_id=2770&sid=1&slc\\_lang=en](https://jrsm.khu.ac.ir/browse.php?a_id=2770&sid=1&slc_lang=en)
- Zhang, X., et al. (2021). Longitudinal association between physical activity and health-related quality of life among community-dwelling older adults: a longitudinal study of Urban Health Centers Europe (UHCE). *BMC Geriatrics*, 21(1), 521. <https://doi.org/10.1186/s12877-021-02452-y>

